



Michael H. Metcalf, M.D.

ROSENBERG COOLEY METCALF
THE ORTHOPEDIC CLINIC AT PARK CITY
www.rcmclinic.com

Rehabilitation:

Biceps Tenodesis, Subacromial Decompression, & Distal Clavicle Resection

Phase 1

Sling

1) Full time for weeks 0-6, abduction pillow is optional, trough is required

Weeks 0- 6

1) **NO ACTIVE BICEPS** (*Includes Elbow Flexion & Forearm Supination*)

2) Begin active shoulder motion in all planes

3) Hand and wrist active motion

4) Initiate independent home motion with caregiver

Home exercise program - Consider pendulums, wand exercise for external rotation

5) Active scapular motion (shrugs, retraction, protraction)

6) **NO PRE's/Strengthening**

Phase 2

Weeks 7- 11

1) Begin Active Biceps

2) Continue to advance shoulder to full active motion (all planes)

3) Continue facilitating scapular stability

4) Emphasize continued independent home stretching to maintain motion

5) May begin low-level cuff strengthening after full AROM has been obtained

6) **NO RESISTED BICEPS**

Phase 3

Week 12

1) Begin PRE's for biceps, advance cuff and scapular resistance as tolerated

2) Return to regular activity as tolerated

Maintenance Program

- Daily Stretching (all planes)

- Strengthening 2-3 times / week

Please check our website for updates: <http://www.rcmclinic.com>

